

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
4 March 2004 (04.03.2004)

PCT

(10) International Publication Number
WO 2004/018425 A1

(51) International Patent Classification⁷: C07D 211/58,
453/02, 265/30, 401/12, 405/12, 413/12, A61K 31/4439,
31/444, A61P 29/00

MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number:
PCT/SE2003/001287

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(22) International Filing Date: 19 August 2003 (19.08.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0202483-4 21 August 2002 (21.08.2002) SE

Declaration under Rule 4.17:

— as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES,
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH,
PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN,
TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO
patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG,
ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU,
TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT,
RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

(71) Applicant (*for all designated States except US*): AS-
TRAZENECA AB [SE/SE]; S-151 85 Södertälje (SE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): CUMMING, John
[GB/GB]; AstraZeneca R & D Alderley, Alderley Park,
Macclesfield, Cheshire SK10 4TG (GB). WINTER, Jon
[GB/GB]; AstraZeneca R & D Alderley, Alderley Park,
Macclesfield, Cheshire SK10 4TG (GB).

(74) Agent: GLOBAL INTELLECTUAL PROPERTY; As-
traZeneca AB, S-151 85 Södertälje (SE).

Published:


— with international search report

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

BEST AVAILABLE COPY

(54) Title: N-4-PIPERIDINYL COMPOUNDS AS CCR5 MODULATORS



(57) Abstract: The invention provides a compound of formula (I): [Chemical formula should be inserted here. Please see paper copy.] wherein R₁, R₂, R₃, R₄, R₅, and R₆ are as defined; or a pharmaceutically acceptable salt thereof or a solvate thereof; compositions containing these compounds, processes for preparing them and their use as modulators of chemokine activity (especially CCR5 activity).

WO 2004/018425 A1

This Page Blank (uspto)

N-4-PIPERIDINYL COMPOUNDS AS CCR5 MODULATORS

The present invention relates to heterocyclic derivatives having pharmaceutical activity, to processes for preparing such derivatives, to pharmaceutical compositions comprising such derivatives and to the use of such derivatives as active therapeutic agents.

Pharmaceutically active piperidine derivatives are disclosed in PCT/SE01/01053, EP-A1-1013276, WO00/08013, WO99/38514 and WO99/04794.

Chemokines are chemotactic cytokines that are released by a wide variety of cells to attract macrophages, T cells, eosinophils, basophils and neutrophils to sites of inflammation and also play a rôle in the maturation of cells of the immune system. Chemokines play an important rôle in immune and inflammatory responses in various diseases and disorders, including asthma and allergic diseases, as well as autoimmune pathologies such as rheumatoid arthritis and atherosclerosis. These small secreted molecules are a growing superfamily of 8-14 kDa proteins characterised by a conserved four cysteine motif. The chemokine superfamily can be divided into two main groups exhibiting characteristic structural motifs, the Cys-X-Cys (C-X-C, or α) and Cys-Cys (C-C, or β) families. These are distinguished on the basis of a single amino acid insertion between the NH-proximal pair of cysteine residues and sequence similarity.

The C-X-C chemokines include several potent chemoattractants and activators of neutrophils such as interleukin-8 (IL-8) and neutrophil-activating peptide 2 (NAP-2).

The C-C chemokines include potent chemoattractants of monocytes and lymphocytes but not neutrophils such as human monocyte chemotactic proteins 1-3 (MCP-1, MCP-2 and MCP-3), RANTES (Regulated on Activation, Normal T Expressed and Secreted), eotaxin and the macrophage inflammatory proteins 1 α and 1 β (MIP-1 α and MIP-1 β).

Studies have demonstrated that the actions of the chemokines are mediated by subfamilies of G protein-coupled receptors, among which are the receptors designated CCR1, CCR2, CCR2A, CCR2B, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CCR10, CXCR1, CXCR2, CXCR3 and CXCR4. These receptors represent good targets for drug development since agents which modulate these receptors would be useful in the treatment of disorders and diseases such as those mentioned above.


The CCR5 receptor is expressed on T-lymphocytes, monocytes, macrophages, dendritic cells, microglia and other cell types. These detect and respond to several

chemokines, principally "regulated on activation normal T-cell expressed and secreted" (RANTES), macrophage inflammatory proteins (MIP) MIP-1a and MIP-1b and monocyte chemoattractant protein-2 (MCP-2).

This results in the recruitment of cells of the immune system to sites of disease. In many diseases it is the cells expressing CCR5 which contribute, directly or indirectly, to tissue damage. Consequently, inhibiting the recruitment of these cells is beneficial in a wide range of diseases.

CCR5 is also a co-receptor for HIV-1 and other viruses, allowing these viruses to enter cells. Blocking the receptor with a CCR5 antagonist or inducing receptor internalisation with a CCR5 agonist protects cells from viral infection.

The present invention provides a compound of formula (I):



wherein:

R¹ is C₁₋₆ alkoxy (optionally substituted by C₁₋₄ alkoxy or phenyl), C₃₋₆ alkenyloxy, phenoxy or piperidin-4-yl (1-substituted by C(O)R⁷ or S(O)₂R⁸);

R² is optionally substituted phenyl, optionally substituted heteroaryl or cycloalkyl;

R^{2a}, R⁴ and R^{4a} are, independently, hydrogen or C₁₋₄ alkyl;

R³ and R^{3a} are, independently, hydrogen or C₁₋₄ alkyl or C₁₋₄ alkoxy;

R⁵ is hydrogen, C₁₋₄ alkyl (optionally substituted by halogen, hydroxy, C₁₋₄ alkoxy, C₃₋₇

cycloalkyl, SH, C₁₋₄ alkylthio, cyano or S(O)_q(C₁₋₄ alkyl)), C₃₋₄ alkenyl, C₃₋₄ alkynyl or C₃₋₇ cycloalkyl;

R⁶ is phenyl, heteroaryl, phenylNH, heteroarylNH, phenyl(C₁₋₂)alkyl, heteroaryl(C₁₋₂)alkyl, phenyl(C₁₋₂ alkyl)NH or heteroaryl(C₁₋₂ alkyl)NH;

R⁷ is C₁₋₆ alkyl (optionally substituted by phenyl, heteroaryl, C₁₋₄ alkoxy, or C₁₋₄ alkoxy(C₁₋₄ alkoxy)), C₁₋₆ alkoxy, phenyl, heteroaryl or C₃₋₆ cycloalkyl;

R⁸ is C₁₋₆ alkyl (optionally substituted by phenyl) or phenyl;

wherein the phenyl and heteroaryl rings of any of the foregoing are independently optionally substituted by halo, cyano, nitro, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, S(O)_mC₁₋₄ alkyl,

S(O)₂NR⁹R¹⁰, NHS(O)₂(C₁₋₄ alkyl), NH₂, NH(C₁₋₄ alkyl), N(C₁₋₄ alkyl)₂, NHC(O)NH₂,

$C(O)NH_2$, $C(O)NH(C_{1-4} \text{ alkyl})$, $NHC(O)(C_{1-4} \text{ alkyl})$, CO_2H , $CO_2(C_{1-4} \text{ alkyl})$, $C(O)(C_{1-4} \text{ alkyl})$, CF_3 , CHF_2 , CH_2F , CH_2CF_3 or OCF_3 ;

R^9 and R^{10} are, independently, hydrogen or C_{1-4} alkyl, or together with a nitrogen or oxygen atom, may join to form a 5- or 6-membered ring which is optionally substituted with C_{1-4}

5 $alkyl$, $C(O)H$ or $C(O)(C_{1-4} \text{ alkyl})$;

m , p and q are, independently, 0, 1 or 2;

or a pharmaceutically acceptable salt thereof or a solvate thereof.

Certain compounds of the present invention can exist in different isomeric forms (such as enantiomers, diastereomers, geometric isomers or tautomers). The present invention covers
10 all such isomers and mixtures thereof in all proportions.

Suitable salts include acid addition salts such as a hydrochloride, hydrobromide, phosphate, acetate, fumarate, maleate, tartrate, citrate, oxalate, methanesulphonate or *p*-toluenesulphonate.

The compounds of the invention may exist as solvates (such as hydrates) and the
15 present invention covers all such solvates.

Alkyl groups and moieties preferably contain, unless otherwise specified, 1-6, especially 1-4, carbon atoms. Alkyl groups and moieties are straight or branched chain and are, for example, methyl, ethyl, *n*-propyl, iso-propyl or tert-butyl.

Alkenyl and alkynyl groups and moieties preferably contain, unless otherwise
20 specified, 2-6, especially 2-4, carbon atoms. Alkenyl includes prop-2-en-1-yl, allyl, but-3-en-1-yl, but-1-en-1-yl, 2-methylallyl, 1-methyl-but-3-en-1-yl, 1-methyl-but-1-en-1-yl, pent-2-en-1-yl and hex-1-en-1-yl. Alkynyl includes propargyl, but-3-yn-1-yl, pent-4-yn-1-yl and hex-5-yn-1-yl. Alkenyl and alkynyl groups and moieties are, for example, vinyl, allyl or propargyl.

Cycloalkyl preferably contains, unless otherwise specified, 3-7, especially 3-6, carbon
25 atoms. Cycloalkyl is, for example, cyclopropyl, cyclobutyl or cyclopentyl.

Cycloalkyl fused to a phenyl ring is, for example, benzocyclobuten-1-yl, indan-1-yl or indan-2-yl.

Heteroaryl is an aromatic 5 or 6 membered ring comprising at least one heteroatom selected from the group comprising nitrogen, oxygen and sulphur. Heteroaryl is, for example,
30 pyrrolyl, imidazolyl, pyrazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, thienyl, furyl, quinolinyl, isoquinolinyl, dihydroisoquinolinyl, quinazolinyl, quinoxalinyl,

indolyl, isoindolyl, benzimidazolyl, benzo[b]furyl, benzo[b]thienyl, phthalazinyl, benzthiazolyl or cinnolinyl.

Phenylalkyl is, for example, benzyl, 1-(phenyl)eth-1-yl or 1-(phenyl)eth-2-yl.

Heteroarylalkyl is, for example, pyridinylmethyl, pyrimidinylmethyl or 1-
5 (pyridinyl)eth-2-yl.

The group $S(O)_2NR^9R^{10}$ is, for example, $S(O)_2NH_2$, $S(O)_2NH(C_{1-4} \text{ alkyl})$, $S(O)_2N(C_{1-4} \text{ alkyl})_2$, $S(O)_2(4-C(O)H\text{-piperazin-1-yl})$ or $S(O)_2(4-C(O)CH_3\text{-piperazin-1-yl})$.

Phenyl(C_{1-2} alkyl)NH is, for example, benzylamino. Heteroaryl(C_{1-2} alkyl)NH is, for example, pyridinylCH₂NH, pyrimidinylCH₂NH or pyridinylCH(CH₃)NH.

10 In one aspect the present invention provides a compound of formula (I), wherein R^1 is piperidin-4-yl (1-substituted by $C(O)R^7$ or $S(O)_2R^8$), wherein R^7 and R^8 are as defined above.

In a further aspect R^1 is piperidin-4-yl 1-substituted by $C(O)R^7$, wherein R^7 is as defined above. R^7 is, for example, C_{1-6} alkyl (optionally mono-substituted by phenyl), C_{1-6} alkoxy, phenyl or C_{3-6} cycloalkyl, wherein the phenyl rings are optionally substituted by
15 halogen.

In a still further aspect R^1 is piperidin-4-yl 1-substituted by $S(O)_2R^8$, wherein R^8 is as defined above. R^8 is, for example, phenyl or C_{1-6} alkyl (optionally mono-substituted by phenyl), wherein the phenyl rings are optionally substituted by halogen, $S(O)_2(C_{1-4} \text{ alkyl})$ or $NHC(O)(C_{1-4} \text{ alkyl})$.

20 In another aspect R^1 is C_{1-6} alkoxy (optionally substituted by C_{1-4} alkoxy or phenyl), C_{3-6} alkenyloxy or phenoxy (optionally substituted by halogen).

In yet another aspect of the present invention R^1 is piperidin-4-yl 1-substituted by $C(O)R^7$ {wherein R^7 is C_{1-6} alkyl (optionally mono-substituted by phenyl), C_{1-6} alkoxy, phenyl or C_{3-6} cycloalkyl, wherein the phenyl rings are optionally substituted by halogen} or $S(O)_2R^8$
25 {wherein R^8 is phenyl or C_{1-6} alkyl (optionally mono-substituted by phenyl), wherein the phenyl rings are optionally substituted by halogen, $S(O)_2(C_{1-4} \text{ alkyl})$ or $NHC(O)(C_{1-4} \text{ alkyl})$ }, or R^1 is C_{1-6} alkoxy (optionally substituted by C_{1-4} alkoxy or phenyl), C_{3-6} alkenyloxy or phenoxy (optionally substituted by halogen).

In a further aspect R^2 is phenyl optionally substituted (such as in the ortho or meta
30 position) by halo, C_{1-4} alkyl, C_{1-4} alkoxy, $S(O)_n(C_{1-4} \text{ alkyl})$ (wherein n is 0, 1 or 2), nitro, cyano or CF_3 . Halo is especially fluorine or chlorine. For example R^2 is unsubstituted phenyl, 3-fluorophenyl, 3-chlorophenyl, 4-fluorophenyl or 4- CF_3 -phenyl. R^2 can additionally be 3,5-difluorophenyl.

In a still further aspect R^2 is phenyl optionally substituted (such as in one of both meta positions) by halo (for example fluorine), C_{1-4} alkyl, cyano or CF_3 . For example R^2 is unsubstituted phenyl, 3-fluorophenyl or 3,5-difluorophenyl.

In another aspect R^2 is phenyl optionally substituted in the ortho or meta position by fluorine or chlorine. For example R^2 is unsubstituted phenyl, 3-fluorophenyl or 3-chlorophenyl.

In yet another aspect R^2 is phenyl optionally substituted (such as in one of both meta positions) by halo (for example fluorine). For example R^2 is unsubstituted phenyl, 3-fluorophenyl or 3,5-difluorophenyl.

10 In another aspect R^{2a} is hydrogen.

In yet another aspect R^3 and R^{3a} are both hydrogen.

In a yet further aspect R^4 and R^{4a} are, independently, hydrogen or methyl (for example R^4 is hydrogen and R^{4a} is methyl, or, R^4 and R^{4a} are both hydrogen).

In a still further aspect R^4 is hydrogen or methyl and R^{4a} is hydrogen.

15 In a still further aspect R^{2a} , R^3 , R^{3a} , R^4 and R^{4a} are all hydrogen.

In another aspect R^5 is hydrogen, C_{1-4} alkyl (such as methyl, ethyl or iso-propyl), C_{3-4} alkenyl, C_{3-4} alkynyl, C_{3-7} cycloalkyl or C_{3-7} cycloalkyl(C_{1-4} alkyl). For example R^5 is hydrogen, methyl, ethyl, allyl, propargyl, cyclopropyl or cyclopropyl CH_2 .

In a further aspect R^5 is hydrogen, methyl, ethyl, allyl or cyclopropyl.

20 In still further aspects of the invention R^5 is ethyl.

In another aspect R^6 is phenyl or benzyl; the phenyl rings of these being optionally substituted by one or more of: halo, cyano, nitro, hydroxy, C_{1-4} alkyl, C_{1-4} alkoxy, $S(O)_m C_{1-4}$ alkyl, $S(O)_2 NR^9 R^{10}$, $NHS(O)_2(C_{1-4}$ alkyl), NH_2 , $NH(C_{1-4}$ alkyl), $N(C_{1-4}$ alkyl) $_2$, $NHC(O)NH_2$, $C(O)NH_2$, $C(O)NH(C_{1-4}$ alkyl), $NHC(O)(C_{1-4}$ alkyl), CO_2H , $CO_2(C_{1-4}$ alkyl), $C(O)(C_{1-4}$ alkyl), CF_3 , CHF_2 , CH_2F , CH_2CF_3 or OCF_3 ; wherein m , R^9 and R^{10} are as defined above or below.

In a still further aspect R^6 is benzyl the phenyl ring of which is optionally substituted by one or more of: halo, cyano, nitro, hydroxy, C_{1-4} alkyl, C_{1-4} alkoxy, $S(O)_m C_{1-4}$ alkyl, $S(O)_2 NR^9 R^{10}$, $NHS(O)_2(C_{1-4}$ alkyl), NH_2 , $NH(C_{1-4}$ alkyl), $N(C_{1-4}$ alkyl) $_2$, $NHC(O)NH_2$, $C(O)NH_2$, $C(O)NH(C_{1-4}$ alkyl), $NHC(O)(C_{1-4}$ alkyl), CO_2H , $CO_2(C_{1-4}$ alkyl), $C(O)(C_{1-4}$ alkyl), CF_3 , CHF_2 , CH_2F , CH_2CF_3 or OCF_3 ; wherein m , R^9 and R^{10} are as defined above or below.


30 In another aspect of the invention R^9 and R^{10} are, independently, hydrogen or C_{1-4} alkyl.

In yet another aspect of the invention R^6 is benzyl singly substituted (such as in the 4-position) by $S(O)_2(C_{1-4})alkyl$ (such as $S(O)_2CH_3$) or $S(O)_2NR^9R^{10}$ (R^9 and R^{10} are, independently, hydrogen or C_{1-4} alkyl, or together with a nitrogen or oxygen atom, may join to form a 5- or 6-membered ring which is optionally substituted with C_{1-4} alkyl, $C(O)H$ or $C(O)(C_{1-4} alkyl)$) (such as $S(O)_2NH_2$, $S(O)_2NH(CH_3)$, $S(O)_2N(CH_3)_2$, $S(O)_2(4-C(O)H-piperazin-1-yl)$ or $S(O)_2(4-C(O)CH_3-piperazin-1-yl)$. The 5- or 6-membered ring is, for example, morpholine, thiomorpholine, piperidine, piperazine or pyrrolidine; but is especially piperazine.

In another aspect of the invention R^6 is benzyl singly substituted (such as in the 4-position) by $S(O)_2(C_{1-4})alkyl$ (such as $S(O)_2CH_3$).

In a further aspect the present invention provides a compound of formula (I) wherein R^1 is piperidin-4-yl 1-substituted by $C(O)R^7$ {wherein R^7 is C_{1-6} alkyl (optionally mono-substituted by phenyl), C_{1-6} alkoxy, phenyl or C_{3-6} cycloalkyl, wherein the phenyl rings are optionally substituted by halogen} or $S(O)_2R^8$ {wherein R^8 is phenyl or C_{1-6} alkyl (optionally mono-substituted by phenyl), wherein the phenyl rings are optionally substituted by halogen, $S(O)_2(C_{1-4} alkyl)$ or $NHC(O)(C_{1-4} alkyl)$ }, or R^1 is C_{1-6} alkoxy (optionally substituted by C_{1-4} alkoxy or phenyl), C_{3-6} alkenyloxy or phenoxy (optionally substituted by halogen); R^2 is phenyl or phenyl substituted in one or both meta-positions by halogen (for example 3-fluorophenyl); R^{2a} , R^3 , R^{3a} , R^4 and R^{4a} are all hydrogen; R^5 is ethyl; and, R^6 benzyl wherein the phenyl ring is substituted by $S(O)_2(C_{1-4} alkyl)$ (for example $S(O)_2CH_3$); or a pharmaceutically acceptable salt thereof (such as a salt of a hydro-halogen acid, for example a hydrochloride).


In yet another aspect the present invention provides a compound of formula (Ia):



wherein R^1 , R^5 and R^6 are as defined above, and R^2 is unsubstituted phenyl or 3-fluorophenyl. In a further aspect the invention provides a compound of formula (I) wherein R^1 , R^5 and R^6 are as defined above, and R^2 is unsubstituted phenyl, 3-fluorophenyl or 3,5-difluorophenyl. In


another aspect of the invention the compounds of formula (Ia) have S absolute configuration at the asterisked carbon (that is, the carbon labelled '*').

In a further aspect the present invention provides a compound of formula (Ib):



- 5 having S absolute configuration at *, wherein R^{1a} is piperidin-4-yl 1-substituted by: $C(O)R^7$ or $S(O)_2R^8$, wherein R^7 is as defined above (for example R^7 is C_{1-6} alkyl (optionally mono-substituted by phenyl), C_{1-6} alkoxy, phenyl or C_{3-6} cycloalkyl, wherein the phenyl rings are optionally substituted by halogen) and R^8 is as defined above (for example R^8 is phenyl or C_{1-6} alkyl (optionally mono-substituted by phenyl), wherein the phenyl rings are optionally substituted by halogen, $S(O)_2(C_{1-4}$ alkyl) or $NHC(O)(C_{1-4}$ alkyl)); or a pharmaceutically acceptable salt thereof (such as a salt of a hydro-halogen acid, for example a hydrochloride).
- 10

In a still further aspect the present invention provides a compound of formula (Ic):




- 15 having S absolute configuration at *, wherein R^1 is as defined above (for example R^1 is C_{1-6} alkoxy (optionally substituted by C_{1-4} alkoxy or phenyl), C_{3-6} alkenyloxy or phenoxy (optionally substituted by halogen); and R^{2b} is hydrogen or halogen (for example fluoro).

The following compounds illustrate the invention.

TABLE I

Table I comprises compounds of formula (Ib), all having S absolute configuration at *.




Compound No.	R ^{1a}	LCMS (MH ⁺)
1	C(O)CH ₃	611
2	C(O)OC(CH ₃) ₃	669
3	S(O) ₂ Phenyl	709
4	S(O) ₂ CH(CH ₃) ₂	675
5	S(O) ₂ CH ₃	647
6	S(O) ₂ CH ₂ Phenyl	723
7	S(O) ₂ CH ₂ CH ₃	661
8	S(O) ₂ (4-S(O) ₂ CH ₃ -C ₆ H ₄)	787
9	S(O) ₂ (4-F-C ₆ H ₄)	727
10	S(O) ₂ (4-NHC(O)CH ₃ -C ₆ H ₄)	766
11	C(O)CH(CH ₃) ₂	639
12	C(O)CH ₂ Phenyl	687
13	C(O)CH ₂ O(CH ₂) ₂ OCH ₃	685
14	C(O)Cyclopropyl	637
15	C(O)(4-Cl-C ₆ H ₄)	751
16	C(O)Phenyl	673

TABLE II

Table II comprises compounds of formula (Ic), all having S absolute configuration at


*



Compound No.	R ¹	R ^{2b}	Salt	LCMS (MH ⁺)
1	OC(CH ₃) ₃	H		558
2	OCH ₂ CH(CH ₃) ₂	H	HCl	558
3	O(CH ₂) ₂ OCH ₃	H	HCl	560
4	Phenoxy	H	HCl	578
5	Benzyloxy	H	HCl	592
6	O(4-F-C ₆ H ₄)	H	HCl	596
7	OCH ₂ CCl ₃	H	HCl	634
8	OCH ₂ CH=CH ₂	H	HCl	542
9	O(4-Cl-C ₆ H ₄)	H	HCl	612
10	OC(CH ₃) ₃	F		576

5

A compound of formula (I), (Ia), (Ib) or (Ic) can be prepared by treating a compound of formula (II):




with:

- 10 an acid chloride or chloroformate of formula R¹C(O)Cl, in the presence of a base (such as a tertiary amine, such as N(C₁₋₆ alkyl)₃ where the alkyl groups can be the same, two of the same

or all different, for example triethylamine) and in a suitable solvent (such as a chlorinated hydrocarbon, for example dichloromethane); or


- when R^1 is a 1-substituted piperidin-4-yl, an acid of formula R^1CO_2H in the presence of a suitable coupling agent (such as O-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate [HATU] or bromo-tris-pyrrolidino-phosphonium hexafluorophosphate [PyBrop]) in the presence of a suitable base (such as a tertiary amine, for example diisopropylethylamine) in a suitable solvent (such as *N*-methylpyrrolidinone).

A compound of formula (II) can be prepared by treating a compound of formula (III):




- with trifluoroacetic acid or hydrochloric acid in the presence of methanol, and then basifying to release the free amine form of formula (II).

A compound of formula (III) can be prepared by reductively aminating a compound of formula (IV):




- with a compound of formula (V):



in the presence of a suitable solvent (such as an aliphatic alcohol such as methanol), a suitable organic acid (such as an aliphatic acid, for example acetic acid) and a suitable reducing agent (such as sodium triacetoxyborohydride or sodium cyanoborohydride).

- A compound of formula (II) wherein R^{2a} is hydrogen can be prepared by reductive amination of a compound of formula (VI):



for example by reacting a compound of formula (VI) with hydroxylamine and hydrogenating the product so formed with hydrogen in the presence of a suitable metal catalyst (such as palladium or platinum catalyst, for example palladium on charcoal).


- 5 A compound of formula (VI), wherein R^{4a} is hydrogen, can be prepared by reacting a compound of formula (V) with:

an alkyl halide of formula $R^2C(O)CR^3R^{3a}CHR^4X$ (wherein X is halogen, such as chloro, bromo or iodo) in the presence of a suitable base (such as potassium carbonate) and a suitable solvent (such as acetone); or,

- 10 compounds of formula $R^2C(O)CHR^3R^{3a}$ and R^4CHO in the presence of a suitable acid (such as acetic acid).

A compound of formula (VI), wherein R^{3a} is hydrogen, can be prepared by reacting a compound of formula (V) with an alkene of formula $R^2C(O)CR^3=CR^4R^{4a}$ in a suitable solvent (such as an aliphatic alcohol, for example ethanol) at a temperature in the range -10 to 100°C .

- 15 Alternatively, a compound of formula (I) can be prepared by reacting a compound of formula (VII):




with an acid chloride $R^7C(O)Cl$ or sulfonyl chloride $R^8S(O)_2Cl$ in the presence of a base (such as a tertiary amine, such as $N(C_{1-6} \text{ alkyl})_3$ where the alkyl groups can be the same, two of the same or all different, for example triethylamine) and in a suitable solvent (such as a chlorinated hydrocarbon, for example dichloromethane); or

- 20 with an acid of formula R^7CO_2H in the presence of a suitable coupling agent (such as O-(7-azabenzotriazol-1-yl)- N,N,N',N' -tetramethyluronium hexafluorophosphate [HATU] or bromo-tris-pyrrolidino-phosphonium hexafluorophosphate [PyBrop]) in the presence of a suitable


base (such as a tertiary amine, for example diisopropylethylamine) in a suitable solvent (such as *N*-methylpyrrolidinone).

A compound of formula (VII) can be prepared by deprotecting a compound of formula (VIII):



with a suitable acid (such as hydrochloric acid or trifluoroacetic acid).

A compound of formula (VIII) can be prepared by reacting a compound of formula (IX):



with 1-Boc-piperidine-4-carboxylic acid and a suitable coupling reagent (such as O-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate [HATU] or bromo-tris-pyrrolidino-phosphonium hexafluorophosphate [PyBrop]).

The starting materials for these processes are commercially available, can be prepared by literature methods or can be prepared by adapting literature methods. In a further aspect the invention provides processes for preparing the compounds of formulae (I), (Ia), (Ib) and (Ic). Many of the intermediates in the processes are novel and these are provided as further features of the invention.

The compounds of the invention have activity as pharmaceuticals, in particular as modulators (such as agonists, partial agonists, inverse agonists or antagonists) of chemokine receptor (especially CCR5) activity, and may be used in the treatment of autoimmune, inflammatory, proliferative or hyperproliferative diseases, or immunologically-mediated diseases (including rejection of transplanted organs or tissues and Acquired Immunodeficiency Syndrome (AIDS)).

The compounds of the present invention are also of value in inhibiting the entry of viruses (such as human immunodeficiency virus (HIV)) into target cells and, therefore, are of value in the prevention of infection by viruses (such as HIV), the treatment of infection by viruses (such as HIV) and the prevention and/or treatment of acquired immune deficiency syndrome (AIDS).

According to a further feature of the invention there is provided a compound of the formula (I), (Ia), (Ib) or (Ic) {for example (I), (Ia) or (Ib)}, or a pharmaceutically acceptable salt thereof or a solvate thereof, for use in a method of treatment of a warm blooded animal (such as man) by therapy (including prophylaxis).

The present invention further provides a method of treating a chemokine mediated disease state (especially a CCR5 mediated disease state, such as rheumatoid arthritis) in a warm blooded animal (such as man) suffering from, or at risk of, said disease, which comprises administering to an animal in need of such treatment a therapeutically effective amount of a compound of formula (I), (Ia), (Ib) or (Ic) {for example (I), (Ia) or (Ib)}, or a pharmaceutically acceptable salt thereof or solvate thereof.

The invention also provides a compound of the formula (I), (Ia), (Ib) or (Ic) {for example (I), (Ia) or (Ib)}, or a pharmaceutically acceptable salt thereof or a solvate thereof, for use in therapy (including prophylaxis); for example in the treatment of a chemokine mediated disease state (especially a CCR5 mediated disease state) in a warm blooded animal, such as man, such as in the treatment of rheumatoid arthritis.

In another aspect the present invention provides the use of a compound of the formula (I), (Ia), (Ib) or (Ic) {for example (I), (Ia) or (Ib)}, or a pharmaceutically acceptable salt thereof or a solvate thereof, in the manufacture of a medicament for use in therapy (for example in modulating chemokine receptor activity (especially CCR5 receptor activity (especially in the treatment of rheumatoid arthritis)) in a warm blooded animal, such as man).

The invention further provides the use of a compound of formula (I), (Ia), (Ib) or (Ic) {for example (I), (Ia) or (Ib)}, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for use in the treatment of one or more of the following disease states:

(1) (the respiratory tract) obstructive diseases of airways including: chronic obstructive pulmonary disease (COPD) (such as irreversible COPD); asthma {such as bronchial, allergic, intrinsic, extrinsic or dust asthma, particularly chronic or inveterate asthma (for example late asthma or airways hyper-responsiveness)}; bronchitis {such as eosinophilic bronchitis}; acute,

allergic, atrophic rhinitis or chronic rhinitis including rhinitis caseosa, hypertrophic rhinitis, rhinitis purulenta, rhinitis sicca or rhinitis medicamentosa; membranous rhinitis including croupous, fibrinous or pseudomembranous rhinitis or scrofulous rhinitis; seasonal rhinitis including rhinitis nervosa (hay fever) or vasomotor rhinitis; sarcoidosis; farmer's lung and related diseases; nasal polyposis; fibroid lung or idiopathic interstitial pneumonia;

(2) (bone and joints) arthrides including rheumatic, infectious, autoimmune, seronegative spondyloarthropathies (such as ankylosing spondylitis, psoriatic arthritis or Reiter's disease), Behcet's disease, Sjogren's syndrome or systemic sclerosis;

(3) (skin and eyes) psoriasis, atopic dermatitis, contact dermatitis or other eczematous dermatides, seborrhoetic dermatitis, Lichen planus, Pemphigus, bullous Pemphigus, Epidermolysis bullosa, urticaria, angiodermas, vasculitides erythemas, cutaneous eosinophilias, uveitis, Alopecia areata or vernal conjunctivitis;

(4) (gastrointestinal tract) Coeliac disease, proctitis, eosinophilic gastro-enteritis, mastocytosis, Crohn's disease, ulcerative colitis, irritable bowel disease or food-related allergies which have effects remote from the gut (for example migraine, rhinitis or eczema);

(5) (Allograft rejection) acute and chronic following, for example, transplantation of kidney, heart, liver, lung, bone marrow, skin or cornea; or chronic graft versus host disease; and/or

(6) (other tissues or diseases) Alzheimer's disease, multiple sclerosis, atherosclerosis,

Acquired Immunodeficiency Syndrome (AIDS), Lupus disorders (such as lupus erythematosus or systemic lupus), erythematosus, Hashimoto's thyroiditis, myasthenia gravis, type I diabetes, nephrotic syndrome, eosinophilia fascitis, hyper IgE syndrome, leprosy (such as lepromatous leprosy), Peridontal disease, Sezary syndrome, idiopathic thrombocytopenia purpura or disorders of the menstrual cycle;

in a warm blooded animal, such as man.

In order to use a compound of the invention, or a pharmaceutically acceptable salt thereof or solvate thereof, for the therapeutic treatment of a warm blooded animal, such as man, in particular modulating chemokine receptor (for example CCR5 receptor) activity, said ingredient is normally formulated in accordance with standard pharmaceutical practice as a pharmaceutical composition.

Therefore in another aspect the present invention provides a pharmaceutical composition which comprises a compound of the formula (I), (Ia), (Ib) or (Ic) {for example (I), (Ia) or (Ib)}, or a pharmaceutically acceptable salt thereof or a solvate thereof (active

ingredient), and a pharmaceutically acceptable adjuvant, diluent or carrier. In a further aspect the present invention provides a process for the preparation of said composition which comprises mixing active ingredient with a pharmaceutically acceptable adjuvant, diluent or carrier. Depending on the mode of administration, the pharmaceutical composition will
5 preferably comprise from 0.05 to 99 %w (per cent by weight), more preferably from 0.05 to 80 %w, still more preferably from 0.10 to 70 %w, and even more preferably from 0.10 to 50 %w, of active ingredient, all percentages by weight being based on total composition.

The pharmaceutical compositions of this invention may be administered in standard manner for the disease condition that it is desired to treat, for example by topical (such as to
10 the lung and/or airways or to the skin), oral, rectal or parenteral administration. For these purposes the compounds of this invention may be formulated by means known in the art into the form of, for example, aerosols, dry powder formulations, tablets, capsules, syrups, powders, granules, aqueous or oily solutions or suspensions, (lipid) emulsions, dispersible powders, suppositories, ointments, creams, drops and sterile injectable aqueous or oily
15 solutions or suspensions.

A suitable pharmaceutical composition of this invention is one suitable for oral administration in unit dosage form, for example a tablet or capsule which contains between 0.1mg and 1g of active ingredient.

In another aspect a pharmaceutical composition of the invention is one suitable for
20 intravenous, subcutaneous or intramuscular injection.

Each patient may receive, for example, an intravenous, subcutaneous or intramuscular dose of 0.01mgkg^{-1} to 100mgkg^{-1} of the compound, preferably in the range of 0.1mgkg^{-1} to 20mgkg^{-1} of this invention, the composition being administered 1 to 4 times per day. The intravenous, subcutaneous and intramuscular dose may be given by means of a bolus
25 injection. Alternatively the intravenous dose may be given by continuous infusion over a period of time. Alternatively each patient will receive a daily oral dose which is approximately equivalent to the daily parenteral dose, the composition being administered 1 to 4 times per day.

The following illustrate representative pharmaceutical dosage forms containing the
30 compound of formula (I), (Ia), (Ib) or (Ic) {for example (I), (Ia) or (Ib)}, or a pharmaceutically acceptable salt thereof or a solvent thereof (hereafter Compound X), for therapeutic or prophylactic use in humans:

(a)

Tablet I	mg/tablet
Compound X	100
Lactose Ph.Eur.	179
Croscarmellose sodium	12.0
Polyvinylpyrrolidone	6
Magnesium stearate	3.0

(b)

Tablet II	mg/tablet
Compound X	50
Lactose Ph.Eur.	229
Croscarmellose sodium	12.0
Polyvinylpyrrolidone	6
Magnesium stearate	3.0

5

(c)

Tablet III	mg/tablet
Compound X	1.0
Lactose Ph.Eur.	92
Croscarmellose sodium	4.0
Polyvinylpyrrolidone	2.0
Magnesium stearate	1.0

(d)

Capsule	mg/capsule
Compound X	10
Lactose Ph.Eur.	389
Croscarmellose sodium	100
Magnesium stearate	1.0

(e)

Injection I	(50 mg/ml)
Compound X	5.0% w/v
Isotonic aqueous solution	to 100%

5 Buffers, pharmaceutically-acceptable cosolvents such as polyethylene glycol, polypropylene glycol, glycerol or ethanol or complexing agents such as hydroxy-propyl β -cyclodextrin may be used to aid formulation.

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. The tablets (a)-(c) may be enteric coated by conventional means, for example to provide a coating of cellulose acetate phthalate.

10 The invention further relates to combination therapies or compositions wherein a compound of formula (I), or a pharmaceutically acceptable salt, solvate or a solvate of a salt thereof, or a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt, solvate or a solvate of a salt thereof, is administered concurrently (possibly in the same composition) or sequentially with an agent for the treatment of any one of the above disease states.

15 In particular, for the treatment of the inflammatory diseases rheumatoid arthritis, psoriasis, inflammatory bowel disease, COPD, asthma and allergic rhinitis a compound of the invention can be combined with a TNF- α inhibitor (such as an anti-TNF monoclonal antibodies (such as Remicade, CDP-870 and D.sub2.E.sub7.), or a TNF receptor immunoglobulin molecule (such as Enbrel.reg.)), a non-selective COX-1 / COX-2 inhibitor
20 (such as piroxicam or diclofenac; a propionic acid such as naproxen, flubiprofen, fenoprofen, ketoprofen or ibuprofen; a fenamate such as mefenamic acid, indomethacin, sulindac or apazone; a pyrazolone such as phenylbutazone; or a salicylate such as aspirin), a COX-2 inhibitor (such as meloxicam, celecoxib, rofecoxib, valdecoxib or etoricoxib) low dose methotrexate, lefunomide; ciclesonide; hydroxychloroquine, d-penicillamine or auranofin, or
25 parenteral or oral gold.

The present invention still further relates to the combination of a compound of the invention together with:

- a leukotriene biosynthesis inhibitor, a 5-lipoxygenase (5-LO) inhibitor or a 5-lipoxygenase activating protein (FLAP) antagonist, such as zileuton, ABT-761,

fenleuton, tepoxalin, Abbott-79175, Abbott-85761, an N-(5-substituted)-thiophene-2-alkylsulfonamide, a 2,6-di-tert-butylphenol hydrazones, a methoxytetrahydropyran such as Zeneca ZD-2138, SB-210661, a pyridinyl-substituted 2-cyanonaphthalene compound such as L-739,010; a 2-cyanoquinoline compound such as L-746,530; an indole or quinoline compound such as MK-591, MK-886 or BAY x 1005;

- a receptor antagonist for a leukotriene LTB₄, LTC₄, LTD₄ or LTE₄ selected from the group consisting of a phenothiazin-3-one such as L-651,392; an amidino compound such as CGS-25019c; a benzoxalamine such as ontazolast; a benzenecarboximidamide such as BIIL 284/260; or a compound such as zafirlukast, ablukast, montelukast, pranlukast, verlukast (MK-679), RG-12525, Ro-245913, iralukast (CGP 45715A) or BAY x 7195;
- a PDE4 inhibitor including an inhibitor of the isoform PDE4D;
- an antihistaminic H₁ receptor antagonist such as cetirizine, loratadine, desloratadine, fexofenadine, astemizole, azelastine or chlorpheniramine;
- a gastroprotective H₂ receptor antagonist;
- an α ₁- and α ₂-adrenoceptor agonist vasoconstrictor sympathomimetic agent, such as propylhexedrine, phenylephrine, phenylpropanolamine, pseudoephedrine, naphazoline hydrochloride, oxymetazoline hydrochloride, tetrahydrozoline hydrochloride, xylometazoline hydrochloride or ethylnorepinephrine hydrochloride;
- an anticholinergic agent such as ipratropium bromide, tiotropium bromide, oxitropium bromide, pirenzepine or telenzepine;
- a β ₁- to β ₄-adrenoceptor agonist such as metaproterenol, isoproterenol, isoprenaline, albuterol, salbutamol, formoterol, salmeterol, terbutaline, orciprenaline, bitolterol mesylate or pirbuterol, or a methylxanthanine including theophylline and aminophylline; sodium cromoglycate; or a muscarinic receptor (M1, M2, and M3) antagonist;
- an insulin-like growth factor type I (IGF-1) mimetic;
- an inhaled glucocorticoid with reduced systemic side effects, such as prednisone, prednisolone, flunisolide, triamcinolone acetonide, beclomethasone dipropionate, budesonide, fluticasone propionate or mometasone furoate;
- an inhibitor of a matrix metalloprotease (MMP), such as a stromelysin, a collagenase, or a gelatinase or aggrecanase; such as collagenase-1 (MMP-1), collagenase-2 (MMP-

8), collagenase-3 (MMP-13), stromelysin-1 (MMP-3), stromelysin-2 (MMP-10), and stromelysin-3 (MMP-11) or MMP-12;

- a modulator of chemokine receptor function such as CCR1, CCR2, CCR2A, CCR2B, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CCR10 and CCR11 (for the C-C family); CXCR1, CXCR2, CXCR3, CXCR4 and CXCR5 (for the C-X-C family) and CX₃CR1 for the C-X₃-C family;
- an osteoporosis agent such as roloxifene, droloxifene, lasofoxifene or fosomax;
- an immunosuppressant agent such as FK-506, rapamycin, cyclosporine, azathioprine or methotrexate; or,
- an existing therapeutic agent for the treatment of osteoarthritis, for example a non-steroidal anti-inflammatory agent (hereinafter NSAID's) such as piroxicam or diclofenac, a propionic acid such as naproxen, flubiprofen, fenoprofen, ketoprofen or ibuprofen, a fenamate such as mefenamic acid, indomethacin, sulindac or apazone, a pyrazolone such as phenylbutazone, a salicylate such as aspirin, a COX-2 inhibitor such as celecoxib, valdecoxib, rofecoxib or etoricoxib, an analgesic or intra-articular therapy such as a corticosteroid or a hyaluronic acid such as hyalgan or synvisc, or a P2X₇ receptor antagonist.

The present invention still further relates to the combination of a compound of the invention together with: (i) a tryptase inhibitor; (ii) a platelet activating factor (PAF) antagonist; (iii) an interleukin converting enzyme (ICE) inhibitor; (iv) an IMPDH inhibitor; (v) an adhesion molecule inhibitor including a VLA-4 antagonist; (vi) a cathepsin; (vii) a MAP kinase inhibitor; (viii) a glucose-6 phosphate dehydrogenase inhibitor; (ix) a kinin-B.sub1. - and B.sub2. -receptor antagonist; (x) an anti-gout agent, e.g., colchicine; (xi) a xanthine oxidase inhibitor, e.g., allopurinol; (xii) an uricosuric agent, e.g., probenecid, sulfinpyrazone or benzbromarone; (xiii) a growth hormone secretagogue; (xiv) a transforming growth factor (TGF β); (xv) a platelet-derived growth factor (PDGF); (xvi) a fibroblast growth factor, e.g., basic fibroblast growth factor (bFGF); (xvii) a granulocyte macrophage colony stimulating factor (GM-CSF); (xviii) a capsaicin cream; (xix) a Tachykinin NK.sub1. and NK.sub3. receptor antagonist selected from the group consisting of NKP-608C; SB-233412 (talnetant); and D-4418; (xx) an elastase inhibitors selected from the group consisting of UT-77 and ZD-0892; (xxi) a TNF α converting enzyme inhibitor (TACE); (xxii) an induced nitric

oxide synthase inhibitor (iNOS); or (xxiii) a chemoattractant receptor-homologous molecule expressed on TH2 cells (a CRTH2 antagonist).

The invention will now be illustrated by the following non-limiting examples in which, unless stated otherwise:

- 5 (i) temperatures are given in degrees Celsius (°C); operations were carried out at room or ambient temperature, that is, at a temperature in the range of 18-25°C;
- (ii) organic solutions were dried over anhydrous magnesium sulphate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 Pascals; 4.5-30 mm Hg) with a bath temperature of up to 60°C;
- 10 (iii) chromatography unless otherwise stated means flash chromatography on silica gel; thin layer chromatography (TLC) was carried out on silica gel plates; where a "Bond Elut" column is referred to, this means a column containing 10g or 20g of silica of 40 micron particle size, the silica being contained in a 60ml disposable syringe and supported by a porous disc, obtained from Varian, Harbor City, California, USA under the name "Mega Bond Elut ST".
- 15 (iv) in general, the course of reactions was followed by TLC and reaction times are given for illustration only;
- (v) yields, when given, are for illustration only and are not necessarily those which can be obtained by diligent process development; preparations were repeated if more material was required;
- 20 (vi) when given, ¹H NMR data is quoted and is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard, determined at 300 MHz using perdeuterio DMSO (CD₃SOCD₃) as the solvent unless otherwise stated; coupling constants (J) are given in Hz;
- (vii) chemical symbols have their usual meanings; SI units and symbols are used;
- 25 (viii) solvent ratios are given in percentage by volume;
- (ix) mass spectra (MS) were run with an electron energy of 70 electron volts in the chemical ionisation (APCI) mode using a direct exposure probe; where indicated ionisation was effected by electrospray (ES); where values for m/z are given, generally only ions which indicate the parent mass are reported, and unless otherwise stated the mass ion quoted is the
- 30 positive mass ion - (M+H)⁺;
- (x) LCMS characterisation was performed using a pair of Gilson 306 pumps with Gilson 233 XL sampler and Waters ZMD4000 mass spectrometer. The LC comprised water symmetry

4.6x50 column C18 with 5 micron particle size. The eluents were: A, water with 0.05% formic acid and B, acetonitrile with 0.05% formic acid. The eluent gradient went from 95% A to 95% B in 6 minutes. Where indicated ionisation was effected by electrospray (ES); where values for m/z are given, generally only ions which indicate the parent mass are reported, and unless otherwise stated the mass ion quoted is the positive mass ion - (M+H)⁺ and

(xi) the following abbreviations are used:

	DMSO	dimethyl sulphoxide;
	DMF	N-dimethylformamide;
	DCM	dichloromethane;
10	HATU	O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate;
	HOBT	1-hydroxybenzotriazole hydrate;
	EDAC	1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride
	DIPEA	N,N-diisopropylethylamine
15	EtOH	ethanol; and
	EtOAc	ethyl acetate.

EXAMPLE 1

This Example illustrates the preparation of (S)-1-acetyl-piperidine-4-carboxylic acid
20 [3-(4-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-piperidin-1-yl)-1-phenyl-propyl]-
amide (Compound No.1 of Table I).

To a stirred solution of (S)-N-[1-(3-amino-3-phenyl-propyl)-piperidin-4-yl]-N-ethyl-2-
(4-methanesulfonyl-phenyl)-acetamide dihydrochloride (Method B, 0.22g, 0.42mmol) in
DCM (5mL) was added N,N-diisopropylethylamine (0.75mL), 1-acetyl-piperidine-4-carboxylic
25 acid (100mg, 0.58mmol) and HATU (300mg) and the resulting mixture was stirred at room
temperature for 18h. The mixture was partitioned between water and DCM, the organic phase
was washed with water and brine, dried (MgSO₄) and concentrated. The residue was purified
by silica column chromatography (eluent 5% MeOH/DCM) to yield the title compound which
was characterised by LCMS: 611 (MH⁺).

EXAMPLE 2

This Example illustrates the preparation of (S)-4-[3-(4-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-piperidin-1-yl)-1-phenyl-propylcarbamoyl]-piperidine-1-carboxylic acid *tert*-butyl ester (Compound No.2 of Table I).

5 To a stirred solution of 1-*tert*-butyloxycarbonylpiperidine-4-carboxylic acid (Method F, 3.5g, 15mmol) in DCM (30mL) was added EDAC (2.87g, 15mmol), HOBT (2.03g, 15mmol) and triethylamine (2.1mL, 15mmol) and the resulting mixture was stirred at room temperature for 5 min. To this mixture was added a solution of (S)-N-[1-(3-amino-3-phenyl-propyl)-piperidin-4-yl]-N-ethyl-2-(4-methanesulfonyl-phenyl)-acetamide dihydrochloride
10 (Method B, 5.3g, 10mmol) and triethylamine (2.8mL, 20mmol) in DCM (30mL) and the resulting mixture was stirred at room temperature for 18h. The mixture was evaporated and the residue partitioned between water and DCM, the organic phase was washed with brine, dried (MgSO₄) and concentrated to give the title compound (5.9g, 88%).

¹H NMR: 1.0 and 1.1 (t, 3H), 1.4 (s, 9H), 1.5-1.7 (m, 10H), 1.9 (m, 2H), 2.3 (m, 4H),
15 2.8 (m, 4H), 3.2 (s, 3H), 3.2 and 3.3 (q, 2H), 3.6 and 4.1 (m, 1H), 3.8 and 3.85 (s, 2H), 3.9 (m, 1H), 4.8 (m, 1H), 7.2 (m, 1H), 7.3 (m, 4H), 7.5 (d, 2H), 7.85 (d, 2H), 8.3 (m, 1H).

LCMS: 669 (MH⁺).

EXAMPLE 3

20 This Example illustrates the preparation of (S)-1-benzenesulfonyl-piperidine-4-carboxylic acid [3-(4-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-piperidin-1-yl)-1-phenyl-propyl]-amide (Compound No.3 of Table I).

To a stirred solution of (S)-piperidine-4-carboxylic acid [3-(4-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-piperidin-1-yl)-1-phenyl-propyl]-amide (Method E,
25 248mg, 0.45mmol) and triethylamine (0.14mL, 1.0mmol) in DCM (2mL) under argon was added a solution of benzenesulfonyl chloride (89mg, 0.5mmol) in DCM (3mL) and the resulting mixture was stirred at room temperature for 18h. The reaction mixture was partitioned between DCM and water and the organic phase was evaporated. The residue was purified by Bond Elut (gradient elution, ethyl acetate to 1:1 methanol/ethyl acetate) giving the
30 title compound (284mg, 83%).

¹H NMR: 1.0 and 1.1 (t, 3H), 1.5-1.7 (m, 10H), 1.9 (m, 2H), 2.2 (m, 4H), 2.3 (m, 2H),
2.8 (m, 2H), 3.2 (s, 3H), 3.2 and 3.3 (q, 2H), 3.6 and 4.1 (m, 1H), 3.6 (m, 1H), 3.8 and 3.9 (s,

2H), 4.8 (m, 1H), 7.2 (m, 6H), 7.5 (m, 2H), 7.6 (m, 2H), 7.7 (m, 2H), 7.85 (d, 2H), 8.2 (m, 1H).

LCMS: 709 (MH⁺).

5 The procedure described in Example 3 can be repeated using different sulfonyl chlorides (such as methanesulfonyl chloride, ethanesulfonyl chloride, 4-methanesulfonylbenzenesulfonyl chloride, 4-fluorobenzenesulfonyl chloride etc.) in place of benzenesulfonyl chloride.

10 EXAMPLE 4

This Example illustrates the preparation of (S)-1-isobutyryl-piperidine-4-carboxylic acid [3-(4-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-piperidin-1-yl)-1-phenyl-propyl]-amide (Compound No.11 of Table I).

To a stirred solution of isobutyryl chloride (54mg, 0.50mmol) in DCM (1mL) was
15 added a solution of (S)-piperidine-4-carboxylic acid [3-(4-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-piperidin-1-yl)-1-phenyl-propyl]-amide (Method E, 193mg, 0.34mmol) and triethylamine (93μL, 0.68mmol) in DCM (2mL) and the resulting mixture was stirred at room temperature for 18h. The mixture was partitioned between DCM and 2N sodium hydroxide solution and the organic phase was evaporated. The residue was purified by
20 Bond Elut (gradient elution, ethyl acetate to 1:1 methanol/ethyl acetate) giving the title compound (142mg).

¹H NMR: 0.95 (d, 6H), 1.0 and 1.1 (t, 3H), 1.25-1.55 (m, 4H), 1.6-2.0 (m, 8H), 2.2 (m, 2H), 2.45 (m, 2H), 2.8 (m, 2H), 3.0 (m, 1H), 3.2 (s, 3H), 3.2 and 3.35 (q, 2H), 3.3 (m, 1H), 3.6 and 4.1 (m, 1H), 3.8 and 3.9 (s, 2H), 3.95 (m, 1H), 4.4 (m, 1H), 4.8 (m, 1H), 7.2 (m, 1H), 7.3
25 (m, 4H), 7.5 (m, 2H), 7.85 (d, 2H), 8.3 (m, 1H).

LCMS: 639 (MH⁺).

The procedure described in Example 4 can be repeated using different acid chlorides (such as phenylacetyl chloride, benzoyl chloride, cyclopropanecarbonyl chloride, 4-chlorobenzoyl chloride etc.) in place of isobutyryl chloride.
30

EXAMPLE 5

This Example illustrates the preparation of (*S*)-[3-(4-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-piperidin-1-yl)-1-phenyl-propyl]-carbamic acid *tert*-butyl ester (Compound No.1 of Table II).

5 To a stirred solution of (*S*)-(3-oxo-1-phenyl-propyl)-carbamic acid *tert*-butyl ester (Method C, 25g, 100 mmol), *N*-ethyl-2-(4-methanesulfonyl-phenyl)-*N*-piperidin-4-yl-acetamide (Method A, 36g, 110 mmol) and glacial acetic acid (6.0mL) in a 1:1 mixture of DCM and methanol (500mL) was added portionwise sodium triacetoxyborohydride (25g, 120 mmol) at ambient temperature. The reaction mixture was stirred for a further 12 hours, then
10 2N sodium hydroxide solution (500mL) was cautiously added during 30 minutes and the resulting mixture extracted into DCM (500mL). The crude product was purified by chromatography on silica eluting with a 9:1 mixture of ethyl acetate and methanol to give the title compound as a colourless gum, which slowly solidified (40g). A portion was crystallised from ethyl acetate to give a white solid, m.p. 115-116°C.

15 ¹H NMR: 1.0 and 1.1 (t, 3H), 1.35 (s, 9H), 1.5 (m, 2H), 1.7 (m, 4H), 1.9 (m, 2H), 2.2 (t, 2H), 2.8 (m, 2H), 3.15 (s, 3H), 3.2 and 3.3 (q, 2H), 3.6 and 4.1 (m, 1H), 3.8 and 3.85 (s, 2H), 4.5 (m, 1H), 7.2 (m, 5H), 7.4 (br d, 1H), 7.5 (d, 2H), 7.8 (d, 2H).

LCMS: 558 (MH+).

EXAMPLE 6

20 This Example illustrates the preparation of (*S*)-[3-(4-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-piperidin-1-yl)-1-phenyl-propyl]-carbamic acid 1',1',1'-trichloroethoxy ester hydrochloride (Compound No.7 of Table II).

To a stirred solution of (*S*)-*N*-[1-(3-Amino-3-phenyl-propyl)-piperidin-4-yl]-*N*-ethyl-2-(4-methanesulfonyl-phenyl)-acetamide dihydrochloride (Method B, 0.16g, 3 mmol) in DCM
25 (10 mL) was added triethylamine (0.13 mL, 9 mmol) and the reaction was cooled to 0 °C under an atmosphere of argon. Trichloroethyl chloroformate (65 mg, 0.3 mmol) was then added in one portion, and the reaction mixtures were stirred overnight. The mixture was then filtered through a diatomaceous earth cartridge, which had been preloaded with saturated
30 sodium bicarbonate solution, followed by a second cartridge loaded with 1N HCl. The crude product was purified by Bond Elut (dichloromethane then 5% methanol in DCM) and the resulting product was isolated as an HCl salt by addition of 1N HCl in diethyl ether followed by trituration to give the title compound as a white solid (130 mg, 65%).

¹H NMR: 1.3 (br s, 3H), 1.9 (br, 2H), 2.4 (br m, 1H), 2.6 (br m, 3H), 2.8 (br m, 2H), 3.0 (br m, 1H), 3.1 (m, 4H), 3.4 (br m, 2H), 3.6 (br m, 2H), 3.8 (s, 2H), 4.7 (m, 4H), 6.2 (m, 1H), 7.3 (m, 5H), 7.4 (d, 2H), 7.9 (d, 2H), 12.5 (br s, 1H).

LCMS: 634 (MH+).

The procedure described in Example 6 can be repeated using different chloroformates (such as phenyl chloroformate, benzyl chloroformate, methoxyethyl chloroformate, 4-fluorophenyl chloroformate etc.) in place of trichloroethyl chloroformate.

EXAMPLE 7

This Example illustrates the preparation of (S)-[3-(4-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-piperidin-1-yl)-1-(3-fluoro-phenyl)-propyl]-carbamic acid *tert*-butyl ester (Compound No.11 of Table II).

To a solution of (S)-[1-(3-fluoro-phenyl)-3-oxo-propyl]-carbamic acid *tert*-butyl ester (Method D, 0.85g, 3.12mmol) in DCM (70mL) and *N*-ethyl-2-(4-methanesulfonyl-phenyl)-*N*-piperidin-4-yl-acetamide (Method A, 1.19g, 3.67mmol) was added glacial acetic acid (one drop) and the resulting mixture was stirred at room temperature for 1h. Sodium triacetoxymethylborohydride (1.4g, 6.4mmol) was added and the resulting mixture was stirred at room temperature for 18 h. The reaction mixture was quenched with water and the organic phase was washed with sodium hydrogen carbonated solution (saturated aqueous) and water, dried (MgSO₄) and concentrated. The crude product was purified by Bond Elut (ethyl acetate then 8% methanol in ethyl acetate) to give the title compound as a solid (1.00g, 55%).

¹H NMR: 1.0 and 1.1 (t, 3H), 1.35 (s, 9H), 1.5 (m, 2H), 1.7 (m, 4H), 1.9 (m, 2H), 2.2 (t, 2H), 2.8 (m, 2H), 3.2 (s, 3H), 3.2 and 3.3 (q, 2H), 3.6 and 4.1 (m, 1H), 3.8 and 3.85 (s, 2H), 4.5 (m, 1H), 7.05 (m, 1H), 7.1 (m, 2H), 7.35 (dd, 1H), 7.5 (br d, 1H), 7.5 (d, 2H), 7.85 (d, 2H).

LCMS: 576 (MH+).


Starting materials are commercially available, have been described in the literature or can be prepared by adaptation of literature methods. Examples of literature methods include: P. Richter, Ch. Garbe and G. Wagner, *E. Ger. Pharmazie*, 1974, 29(4), 256-262; C. Oniscu, D. Nicoara and G. Funieru, "4-(Ureidosulfonyl)phenylacetic acid and its ureide", RO79-

966646, (Romanian document); and M. A. Zahran, M. M. Ali, Y. A. Mohammed and A. A. Shehata, *Int. J. Chem.*, 1993, 4(3), 61.

Method A

5 *N*-Ethyl-2-(4-methanesulfonyl-phenyl)-*N*-piperidin-4-yl-acetamide

Step 1: Preparation of (1-benzyl-piperidin-4-yl)-ethyl-amine dihydrochloride




To a solution of 1-phenylmethyl-4-piperidone (25.0 g, 132 mmol) in THF (250 mL) was added ethylamine hydrochloride (12.0 g, 147 mmol) and methanol (50 mL) and the
10 resulting mixture stirred at room temperature for 10 min. Sodium triacetoxyborohydride (40 g, 189 mmol) was added portionwise and the resulting mixture stirred at room temperature for 1 h. 2M Sodium hydroxide solution (250 mL) was added and the resulting mixture extracted with diethyl ether. The organic extracts were dried (K_2CO_3) and evaporated to give 1-phenylmethyl-4-ethylaminopiperidine as an oil. This was dissolved in ethanol (500 mL) and
15 concentrated hydrochloric acid (20 mL) was added. The resulting crystals were collected, washed with diethyl ether and dried giving the sub-titled compound as a solid (38 g).

1H NMR: ($CDCl_3$): 1.10 (t, 3H), 1.40 (m, 2H), 1.83 (m, 2H), 2.02 (m, 2H), 2.65 (q, 2H), 2.85 (m, 2H), 3.50 (s, 2H), 3.75 (m, 1H), 7.2 - 7.4 (m, 5H).

MS: 219 (MH $^+$).

20

Step 2: Preparation of *N*-(1-benzyl-piperidin-4-yl)-*N*-ethyl-2-(4-methanesulfonyl-phenyl)-acetamide



To a solution of (1-benzyl-piperidin-4-yl)-ethyl-amine dihydrochloride (32.0g, 110mmol) in DCM (500mL) was added *N,N*-diisopropylethylamine (60mL) with stirring to
25 ensure complete dissolution. 4-Methanesulfonylphenylacetic acid (25.0g, 117mmol), 4-Dimethylaminopyridine (4-DMAP) (2.0g) and dicyclohexylcarbodiimide (DCCl) (25.0g, 121


mmol) were added and the resulting mixture was stirred at room temperature for 20 h. The precipitate was removed by filtration and the resulting solution was washed successively with 2N aqueous HCl, water and 1N aqueous NaOH, dried (MgSO₄) and evaporated. The residue was purified by silica gel chromatography (eluent 10% MeOH/ethyl acetate) to afford the sub-
5 titled compound (35 g, 76%).

¹H NMR: 1.00 and 1.14 (t, 3H), 1.45 and 1.70 (m, 2H), 1.95 (br m, 2H), 2.80 (br m, 2H), 3.18 (s, 3H), 3.20 and 3.33 (q, 2H), 3.45 (s, 2H), 3.80 and 3.87 (s, 2H), 3.70 and 4.10 (m, 1H), 7.2 - 7.3 (m, 5H), 7.48 (m, 2H), 7.82 (m, 2H).

MS: 415 (MH⁺).

10

Step 3: Preparation of title compound




To a solution of *N*-(1-benzyl-piperidin-4-yl)-*N*-ethyl-2-(4-methanesulfonyl-phenyl)-acetamide (34g, 82mmol) in ethanol (600mL) was added ammonium formate (40g). The
15 mixture was purged with argon and 30% Pd on carbon (4.2g) was added. The resulting mixture was stirred at reflux for 4 h, then allowed to cool and filtered through diatomaceous earth. The filtrate was evaporated to give a thick oil which solidified on standing to yield the titled compound (24.9 g, 94%).

¹H NMR: 1.02 and 1.15 (t, 3H), 1.4 -1.6 (br m, 4H), 2.45 (m, 2H), 2.93 (br m, 2H),
20 3.18 (s, 3H), 3.20 and 3.32 (q, 2H), 3.72 and 4.18 (m, 1H), 3.80 and 3.87 (s, 2H), 7.50 (m, 2H), 7.85 (m, 2H).

MS: 325 (MH⁺).

Method B

25 Preparation of (*S*)-*N*-[1-(3-Amino-3-phenyl-propyl)-piperidin-4-yl]-*N*-ethyl-2-(4-methanesulfonyl-phenyl)-acetamide dihydrochloride



(S)-[3-(4-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-piperidin-1-yl)-1-phenyl-propyl]-carbamic acid *tert*-butyl ester (Example 5, 20g) was suspended in DCM (200mL) and a solution of hydrochloric acid in methanol (2M, 50mL) was added. The resulting mixture was stirred for 1h then evaporated yielding the title compound as a white solid (19g).


¹H NMR (d6 DMSO at 373K): 1.1 (t, 3H), 1.5 (m, 2H), 1.9 (m, 2H), 2.0 (m, 1H), 2.3 (m, 2H), 3.0 (m, 1H), 3.2 (m, 4H), 3.3 (q, 2H), 3.9 (s, 2H), 4.0 (m, 1H), 4.4 (m, 1H), 7.4 (m, 3H), 7.5 (m, 4H), 7.9 (m, 2H).

MS: 458.

Method C

(S)-(3-Oxo-1-phenyl-propyl)-carbamic acid *tert*-butyl ester

Step 1: Preparation of (S)-*N*-methyl-*N*-methoxy-3-phenyl-3-Bocaminopropionamide




To a solution of (S)-3-phenyl-3-Bocaminopropanoic acid (available from PepTech Corp. of Cambridge, Massachusetts, USA; 4.97g, 18.7mmol) in DCM (100mL) was added DIPEA (14.8mL, 84.8mmol) and *N,O*-dimethylhydroxylamine hydrochloride (2.21g, 22.7mmol) followed by HATU (8.44g, 84.8mmol). The resulting mixture was stirred at room temperature for 18h, diluted with DCM, washed with 2M aqueous sodium hydroxide and water. The organic phase was dried (Na₂SO₄) and concentrated. The residue was purified by silica column chromatography (eluting with isohexane then 3:1 ethyl acetate to isohexane) giving the sub-title compound as a colourless oil (5.58g, 97%).

^1H NMR (CDCl_3): 1.40 (s, 9H), 2.83 (dd, 1H), 3.01 (m, 1H), 3.08 (s, 3H), 3.52 (s, 3H), 5.10 (m, 1H), 7.28 (m, 5H).

MS: 309.

5 Step 2: Preparation of title compound




To a solution of (*S*)-*N*-methyl-*N*-methoxy-3-phenyl-3-Bocaminopropionamide (5.52g, 17.9mmol) in toluene (180mL) at -20°C was added sodium bis(2-methoxyethoxy)aluminium hydride (65% solution in toluene, 35.8mmol) dropwise. The resulting mixture was stirred at -15°C for 1h. The mixture was washed with saturated aqueous sodium dihydrogen phosphate solution (250mL). The organic phase was dried (Na_2SO_4) and concentrated to give the title compound (5g).

^1H NMR: 1.4 (s, 9H), 2.8 (m, 2H), 5.1 (m, 1H), 7.3 (m, 5H), 8.6 (m, 1H), 9.6 (t, 1H).

15 Method D


(*S*)-[1-(3-Fluoro-phenyl)-3-oxo-propyl]-carbamic acid *tert*-butyl ester

Step 1: Preparation of *trans*-3-fluorocinnamic acid *tert*-butyl ester



To a stirred solution of *trans*-3-fluorocinnamic acid (4.34g, 26.1mmol) in toluene (40mL) at 110°C was added *N,N*-dimethylformamide di-*tert*-butyl acetal (25mL, 104mmol) dropwise over 30 min. The resulting mixture was stirred at reflux for a further 4h. The mixture was then cooled to room temperature and washed with water (50mL), saturated aqueous sodium hydrogen carbonate solution (2 x 100mL) and brine (100mL), dried (MgSO_4) and evaporated. The crude product was purified by Bond Elut (isohexane then 2% ethyl acetate in isohexane) to give the sub-title compound as a liquid (3.7g, 64%).


Step 2: Preparation of (3*S*,1'*R*)-3-[benzyl-(1-phenyl-ethyl)-amino]-3-(3-fluoro-phenyl)-propionic acid *tert*-butyl ester



To a stirred solution of (*R*)-(+)-*N*-benzyl- α -methylbenzylamine (4.0mL, 19mmol) in THF (20mL) at -78°C was added *n*-butyl lithium (1.6M in hexanes, 12.5mL, 20mmol) and the resulting mixture was allowed to warm to room temperature over 10 min. before recooling to -78°C . A solution of *trans*-3-fluorocinnamic acid *tert*-butyl ester (3.74g, 16.8mmol) in THF (20mL) was added and the resulting mixture was stirred at -78°C for 2h then quenched by the addition of saturated aqueous ammonium chloride solution (25mL). After warming to room temperature the organic phase was washed with water (2 x 50mL) and brine, dried (MgSO_4) and evaporated. The crude product was purified by Bond Elut (isohexane then 2% ethyl acetate in isohexane) to give the sub-title compound as a gum (5.85g, 80%).

^1H NMR (400MHz, CDCl_3): 1.23 (s, 9H), 1.27 (d, 3H), 2.48 (m, 2H), 3.67 (s, 2H), 3.97 (q, 1H), 4.40 (dd, 1H), 6.93 (ddd, 1H), 7.1-7.4 (m, 13H).

Step 3: Preparation of 3-*tert*-butoxycarbonylamino-3-(3-fluoro-phenyl)-propionic acid *tert*-butyl ester




A stirred mixture of (3*S*,1'*R*)-3-[benzyl-(1-phenyl-ethyl)-amino]-3-(3-fluoro-phenyl)-propionic acid *tert*-butyl ester (5.39g, 12.4mmol), di-*tert*-butyl dicarbonate (2.98g, 13.7mmol) and 20% palladium hydroxide on carbon (0.59g) in ethanol (100mL) was hydrogenated at 5 Bar at room temperature for 24h. The catalyst was removed by filtration through a pad of

Celite® washing through with ethanol. The filtrate was evaporated to give an oil which was partitioned between ethyl acetate and saturated aqueous sodium hydrogen carbonate solution. The organic phase was dried (MgSO₄) and evaporated. The crude product was purified by Bond Elut (eluting with isohexane then 5% ethyl acetate in isohexane) to give the sub-title compound as an oil (3.63g, 86%).

¹H NMR: 1.33 (s, 18H), 2.63 (m, 2H), 4.90 (m, 1H), 7.06 (ddd, 1H), 7.24 (m, 2H), 7.37 (dd, 1H), 7.50 (br d, 1H).


Step 4: Preparation of (S)-[1-(3-fluoro-phenyl)-3-hydroxy-propyl]-carbamic acid *tert*-butyl ester



To a stirred, ice-cooled solution of 3-*tert*-butoxycarbonylamino-3-(3-fluoro-phenyl)-propionic acid *tert*-butyl ester (2.46g, 7.25mmol) in THF (35mL) was added lithium aluminium hydride (1M in THF, 7.50mL, 7.50mmol) dropwise over 20min. The resulting mixture was stirred with warming to room temperature for 2h. The reaction was quenched with water (0.275mL) then 15% aqueous sodium hydroxide (0.275mL) and more water (0.825mL) were added with stirring. The resultant precipitate was removed by filtration washing with THF, and the filtrate was dried (MgSO₄) and evaporated. The crude product was purified by Bond Elut (gradient elution, isohexane to 30% ethyl acetate in isohexane) to give the sub-title compound as an oil (1.26g, 65%).

¹H NMR: 1.4 (s, 9H), 1.75 (m, 1H), 1.85 (m, 1H), 3.3 (m, 1H), 3.4 (m, 1H), 4.5 (dd, 1H), 4.65 (br m, 1H), 7.1 (m + br s, 3H), 7.35 (m, 2H).

Step 5: Preparation of title compound




To a solution of (*S*)-[1-(3-fluorophenyl)-3-hydroxy-propyl]-carbamic acid *tert*-butyl ester (0.85g, 3.2mmol) in DCM (70mL) under argon was added Dess-Martin periodinane (1.48g, 3.5mmol) and the resulting mixture was stirred at room temperature for 2h before the addition of 2M aqueous sodium hydroxide (50mL). The organic layer was dried (MgSO₄) and evaporated to give the title compound (quantitative).

¹H NMR: 1.4 (s, 9H), 2.8 (m, 2H), 5.1 (m, 1H), 7.05 (ddd, 1H), 7.15 (m, 2H), 7.35 (m, 1H), 7.5 (br d, 1H), 9.6 (s, 1H).

10 Method E

(*S*)-Piperidine-4-carboxylic acid [3-(4-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-piperidin-1-yl)-1-phenyl-propyl]-amide



To a stirred solution of (*S*)-4-[3-(4-{ethyl-[2-(4-methanesulfonyl-phenyl)-acetyl]-amino}-piperidin-1-yl)-1-phenyl-propyl]carbamoyl-piperidine-1-carboxylic acid *tert*-butyl ester (Example 2, 5.85g, 8.73mmol) in DCM (45mL) was added trifluoroacetic acid (15mL) and the resulting mixture was stirred at room temperature for 2h. The mixture was evaporated and the residue partitioned between DCM and 2N sodium hydroxide solution. The organic phase was washed with brine, dried (MgSO₄) and evaporated giving the title compound (4.0g, 81%).

¹H NMR: 1.0 and 1.1 (t, 3H), 1.5-1.9 (m, 14H), 2.4 (m, 2H), 2.8 (m, 2H), 3.0 (2H), 3.2 (s, 3H), 3.2 and 3.4 (q, 2H), 3.6 and 4.1 (m, 1H), 3.8 and 3.9 (s, 2H), 4.1 (m, 1H), 4.8 (m, 1H), 7.2 (m, 1H), 7.3 (m, 4H), 7.5 (d, 2H), 7.85 (d, 2H), 8.3 (m, 1H).

LCMS: 569 (MH⁺).

5

Method F

1-*tert*-Butyloxycarbonylpiperidine-4-carboxylic acid

To a stirred solution of isonipecotic acid (4.54g, 35mmol) in 2N aqueous sodium hydroxide (38.5mL, 77mmol) was added portionwise a solution of di-*tert*-butyl dicarbonate (8.4g, 38.5mmol) in THF (20mL). The resulting mixture was stirred at room temperature for 10 3h. The THF was removed by evaporation under reduced pressure and the remaining aqueous mixture was washed with DCM then acidified to pH 4-5 with 1N hydrochloric acid. The resultant precipitate was collected by filtration, washed with water and dried under vacuum at 50°C affording the title compound.

15 ¹H NMR: 1.35 (m, 2H), 1.4 (s, 9H), 1.8 (m, 2H), 2.4 (m, 1H), 2.8 (m, 2H), 3.8 (m, 2H), 12.2 (br s, 1H).


EXAMPLE 8

The ability of compounds to inhibit the binding of RANTES or MIP-1 α was assessed 20 by an *in vitro* radioligand binding assay. Membranes were prepared from Chinese hamster ovary cells which expressed the recombinant human CCR5 receptor. These membranes were incubated with 0.1nM iodinated RANTES or MIP-1 α , scintillation proximity beads and various concentrations of the compounds of the invention in 96-well plates. The amount of iodinated RANTES or MIP-1 α bound to the receptor was determined by scintillation 25 counting. Competition curves were obtained for compounds and the concentration of compound which displaced 50% of bound iodinated RANTES or MIP-1 α was calculated (IC₅₀). The compounds of formula (I) had an IC₅₀ of less than 50 μ M. For example: Compound 1 of Table I has an IC₅₀ of 39nM (that is 39 nanoM); Compound 5 of Table I has an IC₅₀ of 28nM; and, Compound 3 of Table II has an IC₅₀ of 110nM.

30

CLAIMS

1. A compound of formula (I):



5 wherein:

R^1 is C_{1-6} alkoxy (optionally substituted by C_{1-4} alkoxy or phenyl), C_{3-6} alkenyloxy, phenoxy or piperidin-4-yl (1-substituted by $C(O)R^7$ or $S(O)_2R^8$);

R^2 is optionally substituted phenyl, optionally substituted heteroaryl or cycloalkyl;

R^{2a} , R^4 and R^{4a} are, independently, hydrogen or C_{1-4} alkyl;

10 R^3 and R^{3a} are, independently, hydrogen or C_{1-4} alkyl or C_{1-4} alkoxy;

R^5 is hydrogen, C_{1-4} alkyl (optionally substituted by halogen, hydroxy, C_{1-4} alkoxy, C_{3-7} cycloalkyl, SH, C_{1-4} alkylthio, cyano or $S(O)_q(C_{1-4}$ alkyl)), C_{3-4} alkenyl, C_{3-4} alkynyl or C_{3-7} cycloalkyl;

15 R^6 is phenyl, heteroaryl, phenylNH, heteroarylNH, phenyl(C_{1-2})alkyl, heteroaryl(C_{1-2})alkyl, phenyl(C_{1-2} alkyl)NH or heteroaryl(C_{1-2} alkyl)NH;

R^7 is C_{1-6} alkyl (optionally substituted by phenyl, heteroaryl, C_{1-4} alkoxy, or C_{1-4} alkoxy(C_{1-4} alkoxy)), C_{1-6} alkoxy, phenyl, heteroaryl or C_{3-6} cycloalkyl;

R^8 is C_{1-6} alkyl (optionally substituted by phenyl) or phenyl;


20 wherein the phenyl and heteroaryl rings of any of the foregoing are independently optionally substituted by halo, cyano, nitro, hydroxy, C_{1-4} alkyl, C_{1-4} alkoxy, $S(O)_m C_{1-4}$ alkyl, $S(O)_2 NR^9 R^{10}$, $NHS(O)_2(C_{1-4}$ alkyl), NH_2 , $NH(C_{1-4}$ alkyl), $N(C_{1-4}$ alkyl) $_2$, $NHC(O)NH_2$, $C(O)NH_2$, $C(O)NH(C_{1-4}$ alkyl), $NHC(O)(C_{1-4}$ alkyl), CO_2H , $CO_2(C_{1-4}$ alkyl), $C(O)(C_{1-4}$ alkyl), CF_3 , CHF_2 , CH_2F , CH_2CF_3 or OCF_3 ;

25 R^9 and R^{10} are, independently, hydrogen or C_{1-4} alkyl, or together with a nitrogen or oxygen atom, may join to form a 5- or 6-membered ring which is optionally substituted with C_{1-4} alkyl, $C(O)H$ or $C(O)(C_{1-4}$ alkyl);

m , p and q are, independently, 0, 1 or 2;

or a pharmaceutically acceptable salt thereof or a solvate thereof.

2. A compound as claimed in claim 1 wherein R^1 is piperidin-4-yl 1-substituted by $C(O)R^7$ {wherein R^7 is C_{1-6} alkyl (optionally mono-substituted by phenyl), C_{1-6} alkoxy, phenyl or C_{3-6} cycloalkyl, wherein the phenyl rings are optionally substituted by halogen} or $S(O)_2R^8$ {wherein R^8 is phenyl or C_{1-6} alkyl (optionally mono-substituted by phenyl), wherein the phenyl rings are optionally substituted by halogen, $S(O)_2(C_{1-4}$ alkyl) or $NHC(O)(C_{1-4}$ alkyl)}, or R^1 is C_{1-6} alkoxy (optionally substituted by C_{1-4} alkoxy or phenyl), C_{3-6} alkenyloxy or phenoxy (optionally substituted by halogen).
3. A compound as claimed in claim 1 or 2 wherein R^2 is phenyl optionally substituted by halo, C_{1-4} alkyl, C_{1-4} alkoxy, $S(O)_n(C_{1-4}$ alkyl) (wherein n is 0, 1 or 2), nitro, cyano or CF_3 .
4. A compound as claimed in claim 1, 2 or 3 wherein R^{2a} , R^3 , R^{3a} , R^4 and R^{4a} are all hydrogen.
5. A compound as claimed in claim 1, 2, 3 or 4 wherein R^5 is ethyl.
6. A compound as claimed in claim 1, 2, 3, 4 or 5 wherein R^6 is benzyl singly substituted by $S(O)_2(C_{1-4})alkyl$ or $S(O)_2NR^9R^{10}$; wherein R^9 and R^{10} are, independently, hydrogen or C_{1-4} alkyl, or together with a nitrogen or oxygen atom, may join to form a 5- or 6-membered ring which is optionally substituted with C_{1-4} alkyl, $C(O)H$ or $C(O)(C_{1-4}$ alkyl).
7. A process for preparing a compound of formula (I) comprising:
 - a) treating a compound of formula (II):




i. an acid chloride or chloroformate of formula $R^1C(O)Cl$, in the presence of a base and in a suitable solvent; or

- ii. when R^1 is a 1-substituted piperidin-4-yl, an acid of formula R^1CO_2H in the presence of a suitable coupling agent in a suitable solvent;

OR

- b) reacting a compound of formula (VII):



- i. with an acid chloride $R^7C(O)Cl$ or sulfonyl chloride $R^8S(O)_2Cl$ in the presence of a base and in a suitable solvent; or
- ii. with an acid of formula R^7CO_2H in the presence of a suitable coupling agent in the presence of a suitable base in a suitable solvent.

8. A pharmaceutical composition which comprises a compound of the formula (I), or a pharmaceutically acceptable salt thereof or solvate thereof as claimed in claim 1, and a pharmaceutically acceptable adjuvant, diluent or carrier.
9. A compound of the formula (I), or a pharmaceutically acceptable salt thereof or solvate thereof as claimed in claim 1, for use in therapy.
10. A compound of formula (I), or a pharmaceutically acceptable salt thereof or solvate thereof as claimed in claim 1, in the manufacture of a medicament for use in therapy.
11. A method of treating a chemokine mediated disease state in a warm blooded animal suffering from, or at risk of, said disease, which comprises administering to an animal in need of such treatment a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof or solvate thereof as claimed in claim 1.

INTERNATIONAL SEARCH REPORT

International application No. -

PCT/SE 03/01287

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: C07D 211/58, C07D 453/02, C07D 265/30, C07D 401/12, C07D 405/12,
C07D 413/12, A61K 31/4439, A61K 31/444 A61P 29/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: A61K, C07D, A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

BEILSTEIN, STN CAPLUS, EPODOC, WPI DATA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 0187839 A1 (ASTRAZENECA AB), 22 November 2001 (22.11.01), claims 1-7 --	1-11
Y	EP 1013276 A1 (PFIZER INC.), 28 June 2000 (28.06.00), claims 1,4 --	1-11
A	J. Med. Chem., Volume 44, 2001, Akira Naya et al, "Design, Synthesis, and Discovery of a Novel CCR1 Antagonist", pages 1429-1435, table 2 --	1-11

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"B" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

19 November 2003

Date of mailing of the international search report

24-11-2003

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

FERNANDO FARIETA/BS

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 03/01287

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Tetrahedron Letters, Volume 42, 2001, John T. Lai, "Totally hindered phenols. 2,6-Di-t-butyl-4-(1,1-dialkyl-1-acetamide)-phenols and their persistent phenoxy radicals", pages 557-560, table 1 TMP --	1-11
E,A	WO 02070479 A1 (ASTRAZENECA AB), 12 Sept 2002 (12.09.02), claims 1-13 --	1-11
A	WO 0076513 A1 (MECK & CO., INC.), 21 December 2000 (21.12.00), claims 1-24 --	1-11
A	WO 0076973 A1 (MERCK & CO., INC.), 21 December 2000 (21.12.00), claims 1-26 -- -----	1-11

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE03/01287

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 11
because they relate to subject matter not required to be searched by this Authority, namely:
see next sheet
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE03/01287

Claim 11 relates to a method of treatment of the human or animal body by surgery or by therapy/a diagnostic method practised on the human or animal body/Rule 39.1(iv). Nevertheless, a search has been executed for this claim. The search has been based on the alleged effects of the compounds/compositions.

INTERNATIONAL SEARCH REPORT

Information on patent family members

06/09/03

International application No.

PCT/SE 03/01287

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0187839 A1	22/11/01	AU 5898101 A	26/11/01
		BR 0110767 A	11/02/03
		CA 2407258 A	22/11/01
		CN 1441781 T	10/09/03
		CZ 20023777 A	14/05/03
		EP 1289957 A	12/03/03
		GB 0011838 D	00/00/00
		NO 20025430 A	18/12/02
		SK 16152002 A	02/05/03
EP 1013276 A1	28/06/00	AP 200102186 D	00/00/00
		AP 200102187 D	00/00/00
		AU 763644 B	31/07/03
		AU 1290400 A	31/07/00
		AU 1675100 A	31/07/00
		BG 105709 A	28/02/02
		BG 105721 A	28/02/02
		BR 9905977 A	14/01/03
		BR 9916585 A	16/10/01
		BR 9917007 A	30/10/01
		CA 2350073 A	06/07/00
		CA 2350573 A	06/07/00
		CN 1331591 T	16/01/02
		CN 1331691 T	16/01/02
		EE 200100344 A	15/10/02
		EE 200100345 A	16/12/02
		EP 1140085 A	10/10/01
		EP 1140920 A	10/10/01
		GB 9828420 D	00/00/00
		HR 20010468 A	28/02/03
		HR 20010478 A	30/06/02
		HU 0104795 A	29/04/02
		HU 0104910 A	28/10/02
		IL 143510 D	00/00/00
		IL 143512 D	00/00/00
		JP 2000212159 A	02/08/00
		JP 2002533393 T	08/10/02
		JP 2002533461 T	08/10/02
		NO 20013149 A	23/08/01
		NO 20013183 A	08/08/01
		PL 349091 A	01/07/02
		PL 349495 A	29/07/02
		SK 8752001 A	04/02/03
		SK 8762001 A	04/02/03
		TR 200101793 T	00/00/00
		TR 200101867 T	00/00/00
		TR 200200938 T	00/00/00
		US 6586430 B	01/07/03
		WO 0038680 A	06/07/00
		WO 0039125 A	06/07/00
		ZA 200104211 A	14/01/02
		ZA 200104254 A	01/11/02
		GB 9922702 D	00/00/00

INTERNATIONAL SEARCH REPORT

Information on patent family members

06/09/03

International application No.

PCT/SE. 03/01287

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	02070479	A1	12/09/02	AU	2382102 A	01/07/02
				EP	1343945 A	17/09/03
				GB	0105077 D	00/00/00
				GB	0115579 D	00/00/00
				SE	0103797 D	00/00/00
WO	0076513	A1	21/12/00	AU	5473800 A	02/01/01
				US	6506777 B	14/01/03
WO	0076973	A1	21/12/00	AU	5328500 A	02/01/01
				US	6472410 B	29/10/02

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

This Page Blank (uspto)